

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H	A2	(11) International Publication Number: WO 99/58541 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/EP99/03134 (22) International Filing Date: 6 May 1999 (06.05.99) (30) Priority Data: 9809939.3 8 May 1998 (08.05.98) GB 9812743.4 12 June 1998 (12.06.98) GB (71) Applicant (for all designated States except US): BIOCHEMIE S.A. [ES/ES]; Apartado de Correos, 97, E-08400 Granollers (ES). (72) Inventors; and (75) Inventors/Applicants (for US only): BOSCH, Immaculada [ES/ES]; Raval Cortines, 21, E-08500 Vic (ES). CENTEL-LAS, Victor [ES/ES]; Plaza Joan Asina, 8, 4-2-a, E-08440 Cardedeu (ES). DIAGO, José [ES/ES]; Calle Josep Camer, 43, 5-a, E-08400 Granollers (ES). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: IMPROVEMENTS IN MACROLIDE PRODUCTION (57) Abstract A process for the production of azithromycins by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins in the presence of formaldehyde and in the presence of a reducing agent in non-halogenated solvent; and azithromycin which is free from halogenated organic solvent in a stable anhydrous form; or in the form of a solvate with non-halogenated solvent with the exception of water.		


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Improvements in macrolide production

This present invention relates to azithromycins, e.g. a compound of formula



I

5


Azithromycin (9-deoxy-9a-aza-9a-methyl-9a-homoerithromycin A) is a well-known antibacterial agent described e.g. in The Merck Index, 12th Edition (1996), page 157 (946).

10

An azithromycin or azithromycins as used herein includes azithromycin of formula I and a compound of formula I, wherein

- one or more free hydroxyl groups are substituted, e.g. by protecting groups; e.g. conventional hydroxy protecting groups; and/or
- 15 - a methoxy group is replaced by a free hydroxyl group or by a hydroxyl group which is substituted excluding methoxy, e.g. by protecting groups, e.g. conventional hydroxy protecting groups; and/or
- a dimethylamino group is replaced by a substituted amino group excluding dimethylamino, e.g. by a protected amino group, e.g. a conventional protected
- 20 amino group.

Azithromycins, e.g. azithromycin of formula I, may e.g. be produced by methylation e.g. by a method as conventional, of 9-deoxo-9a-aza-9a-homoerythromycins, e.g. 9-deoxo-9a-aza-9a-homoerythromycin, e.g. a compound of formula



5

e.g. via an Eschweiler-Clarke reaction, e.g. in the presence of formaldehyde and in the presence of a reducing agent, e.g. formaldehyde/formic acid, in halogenated solvent, such as CHCl_3 .

10 A 9-deoxo-9a-aza-9a-homoerythromycin or 9-deoxo-9a-aza-9a-homoerythromycins as used herein include a compound of formula II and a compound of formula II, wherein

- one or more free hydroxyl groups are substituted, e.g. by protecting groups; e.g. conventional hydroxy protecting groups; and/or
- a methoxy group is replaced by a free hydroxyl group or by a hydroxyl group which is substituted excluding methoxy, e.g. by protecting groups, e.g. conventional hydroxy protecting groups; and/or
- 15 - a dimethylamino group is replaced by a substituted amino group excluding dimethylamino, e.g. by a protected amino group, e.g. conventional protected amino group.

20 9-deoxo-9a-aza-9a-homoerythromycins are known and obtainable e.g. by a method as conventional.

It was now surprisingly found, that an Eschweiler-Clarke reaction in the production of azithromycins starting from 9-deoxo-9a-aza-9a-homoerythromycins including

methylation and isolation of azithromycins, e.g. a compound of formula I, may be carried out avoiding halogenated solvent.

In one aspect the present invention provides a process for the production of azithromycins, e.g. azithromycin, e.g. of formula I, by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins, e.g. a compound of formula II, in the presence of formaldehyde and in the presence of a reducing agent, e.g. formic acid or hydrogen, comprising that methylation, e.g. and isolation of azithromycins, is carried out in non-halogenated solvent, e.g. selected from water, alcohols, ketones, alkyl esters and ethers.

A process according to the present invention may be carried out as follows:

The nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins may be methylated in the presence of formaldehyde and in the presence of a reducing agent in non-halogenated solvent.

A hydroxyl protecting group in 9-deoxo-9a-aza-9a-homoerythromycins includes e.g. conventional hydroxyl protecting groups, including e.g. protecting groups for the diol function present in positions 11 and 12 of a compound of formula II, such as protecting groups which form a cyclic group together with two hydroxy groups, e.g. a hydrogenorthoborate group.

Non-halogenated solvent includes e.g. water and an organic solvent, the chemical formula of which does not contain halogen atoms. In contrast to that, halogenated solvent includes organic solvent, the chemical formula of which contains at least one halogen atom.

Non-halogenated solvent as used herein includes solvent which is miscible with water and solvent which is able to form a two phase system with water, including, e.g. a solvent of a low dielectric constant, e.g. appropriate ketones, such as dialkyl ketones, e.g. for example di(C₁₋₅)alkylketones, e.g. methyl-(C₁₋₅)alkyl ketones, e.g. acetone and methylisobutyl ketone; di-(C₂₋₅) alkyl ketones, such as diethylketone and diisopropylketone; esters, such as alkanolic acid esters, e.g. (C₁₋₆)alkanoic acid, e.g. acetic acid alkyl, e.g. (C₁₋₈) alkyl, such as (C₁₋₆)alkyl, e.g. (C₁₋₄)alkyl esters, for example n-butyl acetate, isopropyl acetate and ethyl acetate; ethers such as cyclic ethers, e.g. tetrahydrofuran and non-cyclic, e.g. dialkyl ethers, such as di(C₁₋₈)alkyl, e.g. di(C₁₋₄)alkyl ethers, such as methyl(C₁₋₄)alkyl ethers, e.g. t-butyl methyl ether; and aromatic hydrocarbons, such as xylene or toluene; and solvents of higher dielectric constant, e.g. alcohols, such as alkyl alcohols, e.g.(C₁₋₅)alkyl alcohols, for example

methanol, ethanol and isopropanol; and amides, e.g. derived from a carboxylic acid, such as an alkanolic acid, , e.g. (C₁₋₃)alkanoic acid, and including monoalkyl- and dialkylamides; e.g. mono(C₁₋₄)alkyl and di(C₁₋₄)alkylamides, such as N-methylformamide, dimethylacetamide and dimethylformamide.

- 5 Preferred non-halogenated solvents includes solvents, with low toxicity, e.g. pharmaceutically acceptable solvents, for example water; and alcohols such as ethanol, isopropanol and butanols; ketones such as acetone, methyl ethyl ketone and methyl isobutyl ketone; alkyl esters such as formic acid and acetic acid esters, e.g. ethyl acetate, isopropyl acetate and butyl acetate and ethers such as tert-butyl methyl ether
 10 and tetrahydrofuran, more preferably alcohols, esters, e.g. acetates, ketones and water. Non-halogenated solvent includes a solvent system, e.g. mixtures of individual solvents, e.g. as described above.

- A solution of 9-deoxo-9a-aza-9a-homoerythromycins in a non-halogenated organic solvent as used as starting material according to the present invention may contain,
 15 e.g. a small amount of, water, e.g. to facilitate dissolution of 9-deoxo-9a-aza-9a-homoerythromycins, e.g. in case that 9-deoxo-9a-aza-9a-homoerythromycins in the form of a salt are used as a starting material.

If not otherwise defined herein alkyl includes (C₁₋₁₂)alkyl, such as (C₁₋₆)alkyl, for example (C₁₋₄)alkyl. Lower alkyl includes e.g. (C₁₋₄)alkyl.

- 20 Formaldehyde according to the present invention as used herein includes formaldehyde, e.g. in the form of a gas, formaldehyde in aqueous and in non-halogenated solvent, e.g. as described above, solution and a formaldehyde precursor. A formaldehyde precursor includes a compound which may set free formaldehyde in an reaction medium, such as paraformaldehyde or s-trioxane. If a non-halogenated
 25 solvent is water, formaldehyde in a molar excess (in respect with a 9-deoxo-9a-aza-9a-homoerythromycins A used as a starting material) in form of an aqueous solution may be used, e.g. without use of further water.

- A reducing agent according to the present invention as used herein includes conventional reducing agents useful in a reductive alkylation process, such as formic
 30 acid, zinc/HCl, sodium cyanoborohydride, sodium triacetoxyborohydride, sodium borohydride, BH₃-pyridine, formic acid and hydrogen, e.g. in the presence of a, e.g. conventional, hydrogenation catalyst. A preferred reducing agent includes e.g. formic acid and hydrogen in the presence of a hydrogenation catalyst.

- The amounts of formaldehyde and of the reducing agent present in the methylation
 35 reaction according to the present invention are not critical; e.g. a stoichiometric amount or more relating to a 9-deoxo-9a-aza-9a-homoerythromycin A starting

compound may conveniently be present. An appropriate amount includes e.g. 1 to 30, such as 1 to 5, e.g. 1 to 3 equivalents of formaldehyde and of a reducing agent.

Appropriate reaction conditions for the methylation reaction according to the present invention include e.g.

- 5 - a temperature range of ca -10°C up to the reflux temperature of the solvent or solvent system present, such as from 30°C to 80°C, e.g. more than 50°C;
- an appropriate pressure, e.g. atmospheric pressure, and a pressure which is above or below atmospheric pressure; and
- appropriate dilution, e.g. a dilution range between 1 g and 1000 g of a 9-deoxo-10 9a-aza-9a-homoerythromycin A starting compound per liter of non-halogenated solvent.

A compound of formula I or a compound of formula I wherein

- one or more free hydroxyl groups are substituted, e.g. by protecting groups; e.g. conventional hydroxy protecting groups; and/or
- 15 - a methoxy group is replaced by a free hydroxyl group or by a hydroxyl group which is substituted excluding methoxy, e.g. by protecting groups, e.g. conventional hydroxy protecting groups; and/or
- a dimethylamino group is replaced by a substituted amino group excluding dimethylamino, e.g. by a protected amino group, e.g. conventional protected amino 20 group,

- may be formed and may be isolated from the reaction mixture, e.g. as conventional, e.g. by removal of solvent from the reaction mixture, e.g. by concentration such as evaporation, e.g. to dryness or almost dryness, e.g. until crystallisation or precipitation of an azithromycin, e.g. of formula I, occurs; or by extraction, e.g. as a salt, e.g. into an 25 aqueous solution of acidic pH; and precipitation, e.g. as a salt from an aqueous medium; or extraction at basic pH into another non-halogenated solvent, e.g. as defined above, which may be the same or different from that used in methylation; and precipitation or crystallization of an azithromycin, e.g. of formula I.

If a compound of formula I, wherein

- 30 - one or more free hydroxyl groups are substituted, e.g. by protecting groups; e.g. conventional hydroxy protecting groups; and/or
- a methoxy group is replaced by a free hydroxyl group or by a hydroxyl group which is substituted excluding methoxy, e.g. by protecting groups, e.g. conventional hydroxy protecting groups; and/or

- a dimethylamino group is replaced by a substituted amino group excluding dimethylamino, e.g. by a protected amino group, e.g. conventional protected amino group,

is formed, this compound may be converted into a compound of formula I, e.g. using

5 e.g. a method as conventional, e.g.

- de- and/or reprotecting one or more hydroxyl groups; and/or
- producing a methoxy group e.g. by methylation of an hydroxy group; and/or
- producing a dimethylamino group e.g. by de- and or realkylation

to obtain a compound of formula I.

10

In another aspect the present invention provides a process for the production of azithromycin, e.g. of formula I, by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins, e.g. a compound of formula II, in the presence of formaldehyde and in the presence of a reducing agent,

15 e.g. formic acid or hydrogen, and, e.g. if necessary, converting a 9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin A into a compound of formula I, comprising that methylation, e.g. and isolation of a compound of formula I, is carried out in non-halogenated solvent, e.g. selected from water, alcohols, ketones, alkyl esters and ethers.

20

In a process according to the present invention high yields may be obtained, for example, methylation may result in a yield higher than 90% or even more, e.g. higher than 95 %; and reaction times may be short, e.g. shorter than 4 hours, such as 2 hours. A process according to prior art, e.g. in halogenated solvent, may e.g. result in

25 lower yields and e.g. higher reaction times.

9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin A, e.g. of formula I obtained according to the present invention is useful, e.g. for the production of a pharmaceutical composition; and, if necessary may be converted into a suitable form

30 for pharmaceutical administration, e.g. into the form of a, e.g. pharmaceutically acceptable salt, and/or e.g. into the form of a solvate, e.g. a hydrate such as a monohydrate or a dihydrate. A dihydrate of a compound of formula I may be e.g. obtained by a method as conventional, such as by crystallisation of a compound of formula I, e.g. in acetone/water or by crystallisation in a mixture of aqueous

35 tetrahydrofuran and hexane. Azithromycin in the form of a monohydrate may be obtained, e.g. as conventional, e.g. by crystallization of azithromycin in ethanol/water.

A compound of formula I may be obtained in the form of a solvate with non-halogenated solvent.

5 In a further aspect, the present invention provides a process for the production of azithromycin, e.g. of formula I, in the form of a solvate, by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins in the presence of formaldehyde and in the presence of a reducing agent, e.g. formaldehyde or hydrogen, and, if necessary, converting a 9-deoxo-9a-aza-10 9a-methyl-9a-homoerithromycin A obtained into a compound of formula I, and, converting a compound of formula I into e.g. a salt and/or, a solvate thereof, comprising that methylation, e.g. and isolation of azithromycin, is carried out in non-halogenated solvent e.g. selected from water, alcohols, ketones, alkyl esters and ethers.

15 In another aspect the present invention provides a process for the production of a pharmaceutical composition comprising azithromycin in e.g. pharmaceutically acceptable salt form or, free form, e.g. in the form of a solvate, in association with at least one pharmaceutical carrier or diluent wherein azithromycin is obtained by a process according to the present invention.

20

The process according to the present invention may e.g. have advantages as follow:

- Azithromycin, e.g. of formula I, e.g. in the form of a solvate, may be obtained without the use of halogenated solvents
 - Azithromycin, e.g. of formula I, e.g. in the form of a solvate may be obtained in
- 25 high yields and short reaction times.

In another aspect surprisingly a process for the production of azithromycin in non-halogenated organic solvent with the exception of water as a solvent was found which provides anhydrous azithromycin in a stable, e.g. non-hygroscopic, form and practically free from halogenated organic solvent.

30 The term " free from halogenated organic solvent" used herein, includes e.g. traces of halogenated solvents in azithromycin according to the present invention, such as amounts of 60 ppm and below, e.g. 2 to 60 ppm, or even less.

A stable, e.g. non-hygroscopic, anhydrous form of azithromycin includes a compound of formula I which is practically free from halogenated solvent and having and

35 maintaining a water content of (about) 0.1% to 2.0%, such as 0.2% to 0.9% under

normal environment air humidity conditions, e.g. for at least 24 hours, e.g. for several weeks.

In another aspect the present invention provides azithromycin, e.g. of formula I, which is free from halogenated organic solvent in a stable, e.g. non-hygroscopic, anhydrous form, e.g. having, e.g. and maintaining, a water content of 0.2% to 0.9%, e.g. under normal environment air humidity conditions, e.g. for, e.g. at least, 24 hours.

Azithromycin in a stable anhydrous form, which is free from halogenated organic solvent according to the present invention may be produced e.g. by producing a solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent, e.g. drying said solution, and isolating and drying a composition containing azithromycin and non-halogenated organic solvent; e.g. directly by an Eschweiler-Clark reaction starting from 9-deoxo-9a-aza-9a-homoerythromycin, e.g. as described above.

For the production of a solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent as used as starting material according to the present invention azithromycin in any form may be used, e.g. in free base form and in the form of a salt; e.g. in non-solvate form, e.g. anhydrous form, and/or in the form of a solvate. A salt includes e.g. preferably a hydrochloride, dihydrochloride, acetate, sulphate. A solvate includes a hydrate, such as a monohydrate and a dihydrate. If a salt of azithromycin is used as a starting material, a base, including e.g. inorganic bases such as sodium and potassium hydroxide, may be used, e.g. may be added to the solution, to provide azithromycin in free base form, e.g. by adjustment of the pH of the solution to an pH where azithromycin is present in free base form, including e.g. a pH range of 7 to 11. The concentration of azithromycin in a solution of a non-halogenated organic solvent is not critical. E.g. for ecological reasons preferably a saturated or almost saturated solution may be used. The temperature for the production of a solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent includes a temperature at which azithromycin is not degraded including e.g. a temperature range of e.g. ca. 0°C to ca. 30°C, such as 0 to 30°C.

Preferred non-halogenated solvents with the exception of water as a solvent include pharmaceutically acceptable solvents, for example alcohols such as ethanol, isopropanol or butanol; ketones such as acetone, methyl ethyl ketone or methyl isobutyl ketone; alkyl esters such as formate or acetate esters, e.g. ethyl acetate,

isopropyl acetate or butyl acetate and ethers such as tert-butyl methyl ether or tetrahydrofuran.

A solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent may be dried e.g. as conventional, e.g. by addition of a drying agent, such as anhydrous sodium or magnesium sulphate or a molecular sieve.

For the isolation of azithromycin in anhydrous form according to the present invention the organic solvent of a solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent may be removed, e.g. by distillation, such as vacuum distillation. Solvent may be removed partially, e.g. until azithromycin in anhydrous form crystallizes or precipitates or (almost) totally, e.g. to dryness. A composition containing azithromycin in anhydrous form and non-halogenated solvent, e.g. 10% and less, such as ca. 1% to 10%, e.g. 1% to 7%; may be obtained. In some cases, e.g. if the solution of azithromycin in non-halogenated solvents contains acetates, alcohols or ethers, azithromycin in the form of a solvate, e.g. crystalline, with non-halogenated solvent may be obtained.

In another aspect the present invention provides a composition containing azithromycin of formula I and, in an amount of 10% and less, a non-halogenated organic solvent with the exception of water and ethanol as a solvent; and, in a further aspect, azithromycin of formula I in the form of a solvate with non-halogenated organic solvent with the exception of water as a solvent.

A composition containing azithromycin in anhydrous form and non-halogenated solvent with the exception of water as a solvent, e.g. in the form of a solvate with non-halogenated solvent, may be dried to obtain azithromycin in anhydrous, e.g. in non-hygroscopic, form and free from halogenated solvents. Appropriate drying temperatures includes a temperature of ca. 40°C and more, such as 40°C to 65°C, e.g. 55°C to 65°C. An appropriate drying time may be used, including for example more than 6 hours, e.g. overnight or more, e.g. 10 to 20 hours, such as 12 to 16 hours.

Anhydrous azithromycin obtained according to the present invention may be non-hygroscopic, e.g. it may maintain a water content below the stoichiometric amount of the monohydrate of 2.3 %, for example below 2 %, e.g. below 1%, e.g. between 0.2 to 0.9%, such as 0.4 to 0.8% even after several hours, e.g. 24 hours, or several weeks, under normal, e.g. normal air humidity, environmental conditions.

Azithromycin which is free from halogenated organic solvent in a stable anhydrous form according to the present invention may be obtained in high purity of 95% and more, e.g. 95% to ca. 98%.

- 5 In another aspect the present invention provides a process for the production of azithromycin which is free from halogenated organic solvent in a stable anhydrous form, comprising producing a solution of azithromycin in a non-halogenated solvent with the exception of water as a solvent, e.g. and drying a solution of azithromycin in a non-halogenated solvent, and isolating and drying a composition containing
10 azithromycin and non-halogenated solvent with the exception of water as a solvent.

- Azithromycin which is free from halogenated organic solvent in a stable anhydrous form, e.g. and azithromycin in the form of a solvate with non-halogenated solvent with the exception of water as a solvent, according to the present invention is useful
15 for the production of a pharmaceutical composition containing azithromycin. A pharmaceutical composition containing azithromycin which is free from halogenated organic solvent in a stable anhydrous form, e.g. and azithromycin in the form of a solvate with non-halogenated solvent with the exception of water as a solvent, may contain the same concentrations of azithromycin and may be used for the same
20 indications in the same dosages as a known pharmaceutical composition containing azithromycin in the form of a monohydrate or dihydrate, e.g. as is presently on the market.

- In another aspect the present invention provides a pharmaceutical composition,
25 comprising a therapeutically effective amount of azithromycin which is free from halogenated organic solvent in a stable anhydrous form in association with at least one pharmaceutical carrier or diluent; and a pharmaceutical composition, comprising a therapeutically effective amount of azithromycin in the form of a solvate with non-halogenated solvent with the exception
30 of water as a solvent in association with at least one pharmaceutical carrier or diluent.

- The following examples illustrate the invention. All temperatures are in degree Centigrade and are uncorrected. The content of azithromycin in stable anhydrous azithromycin obtainable according to the examples is calculated by HPLC on
35 anhydrous basis. Azithromycin used as starting material is obtained by know methods or by a method according to the present invention.

Examples 1 to 9**General Reaction Procedure**

- 0.55 g of 9-deoxo-9a-aza-9a-homoerythromycin A dissolved in 6 ml of a solvent as specified in TABLE 1 below under "Solvent" are treated at room temperature with 0.063 ml of formic acid (98-100 %) and 0.135 ml of 37-38 % w/w aqueous formaldehyde. In example 9 instead of 6 ml of "Solvent" and instead of 0.135 ml of 37-38 % w/w aqueous formaldehyde, 3.9 ml of 37-38 % w/w aqueous formaldehyde are used. The reaction mixture is heated at a temperature as specified in TABLE 1 below under "Temperature (in °Celsius)". Within the time as specified in TABLE 1 below under "Time (in hours)" a conversion of 9-deoxo-9a-aza-9a-homoerythromycin A into azithromycin as specified in TABLE 1 under "Conversion (in % of starting material)" is determined by HPLC.

TABLE 1

Example	Solvent	Temperature	Time	Conversion
1	iso-propanol	65-70	4	95
2	2-butanol	70	2	not determined
3	iso-propyl acetate	70	2	not determined
4	n-butyl acetate	70	2	not determined
5	ethyl acetate	70	2	not determined
6	acetone	65-70	2	94
7	iso-butyl ketone	65-70	2	73
8	toluene	65-70	2	62
9	water	70	4.5	63

15

Example 10**General Isolation Procedure (solvent soluble in water)**

- A reaction mixture obtained as described in Example 1 is cooled to room temperature and the solvent is removed by evaporation.
- 0.52 g (92% of theory) of azithromycin are obtained.
- Such an isolation procedure may be convenient in case that water soluble solvent is used.

Examples 2a to 5a

- General Isolation Procedure (solvent may form a two phase system with water)**

A reaction mixture as obtained in one of the Examples above as specified in TABLE 2 below under "Reaction Example" is cooled to room temperature and treated with 40 ml of solvent as specified in TABLE 2 below under "Solvent", 40 ml of water and 20% sulfuric acid to obtain a pH of the mixture of ca. 4. The mixture obtained is stirred for ca. 15 minutes and the phases are separated. The aqueous phase obtained is extracted with 25 ml of solvent as specified in TABLE 2 below under "Solvent". The aqueous phase is treated with 20% sodium hydroxide to obtain a pH 9 and the mixture obtained is stirred for ca. 15 minutes and extracted with 25 ml of solvent as specified in TABLE 2 below under "Solvent". The organic phase is dried over anhydrous sodium sulphate and the sodium sulphate is filtrated off. The organic solvent is evaporated off.

Azithromycin in a yield as specified in TABLE 2 below under "Yield (in gram)" in % of theory as specified in TABLE 2 below under "% of theory" is obtained.

15

TABLE 2

Example	Reaction Example	Solvent	Yield	% of theory
2a	2	2-butanol	0.51	90
3a	3	iso-propyl acetate	0.46	82
4a	4	n-butyl acetate	0.50	89
5a	5	ethyl acetate	0.54	96

Example 11-18General procedure for obtaining azithromycin in stable, anhydrous form

Azithromycin is dissolved in a solvent as specified in TABLE 3 below. Anhydrous sodium sulphate is added to 100 ml of a solution of azithromycin in a solvent as specified in TABLE 3 below under "solvent" containing azithromycin in a concentration as specified in TABLE 3 below under "conc (mg/ml)". The solid is filtrated off. The solvent of the filtrate is distilled off, e.g. by evaporation, to dryness. Anhydrous azithromycin in an amount (in gram) as specified in TABLE 3 below under "Yield wet (g)" (containing solvent as specified in TABLE 3 below under "solvent") with a content of azithromycin (in %) calculated on anhydrous basis as specified in TABLE 3 below under "Content wet" is obtained and dried at ca. 60°C for ca. 14 hours. Azithromycin in anhydrous form is obtained containing an amount of azithromycin (in %) calculated on anhydrous basis as specified in TABLE 3 under

"Content dry" below and having a water content (in %) as specified in TABLE 3 below under "Content water".

TABLE 3


Example	Solvent	Conc (mg/ml)	Yield wet (g)	Content wet	Content dry	Content water
11	ethyl acetate	30	3.15	90.5%	97.8	0.53
12	isopropyl acetate	30	3.33	89.9%	97.8%	0.53
13	n-butyl acetate	30	3.28	89.5%	97.8%	0.53
14	acetone	30	3.18	95.5%	96.6	0.75
15	methyl- isobutyl ketone	30	3.22	89%	96.6	0.75
16	iso- propanol	30	3.32	92%	97.7	0.55
17	ethanol	30	3.0	-	96.1	0.56
18	methanol	30	3.61	-	97.4	0.61

5

The water content is maintained under normal environment air humidity conditions, for at least 24 hours, e.g. for several weeks.

Claims


1. A process for the production of azithromycins by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins in the presence of formaldehyde and in the presence of a reducing agent, comprising that methylation is carried out in non-halogenated solvent.
2. A process according to claim 1 in the production of azithromycin of formula



10

comprising converting a 9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin A into a compound of formula I.

3. A process according to any one of claims 1 or 2 wherein a 9-deoxo-9a-aza-9a-homoerythromycin is a compound of formula



II

4. A process according to any one of claims 1 to 3, wherein non-halogenated solvent is selected from water, alcohols, ketones, alkyl esters and ethers.

5

5. Azithromycin of formula I according to claim 2 which is free from halogenated organic solvent in a stable anhydrous form.

6. Azithromycin according to claim 5 having a water content of 0.1% to 2.0%.

10

7. Azithromycin according to any one of claims 5 or 6 maintaining a water content of 0.1% to 2.0% under normal environment air humidity conditions for at least 24 hours.

15

8. A process for the production of azithromycin of formula I according to claim 2 which is free from halogenated organic solvent in a stable anhydrous form, comprising producing a solution of azithromycin in a non-halogenated organic solvent with the exception of water as a solvent, and isolating and drying a composition containing azithromycin and non-halogenated organic solvent with the exception of water as a solvent.

20

9. Azithromycin of formula I according to claim 2 in the form of a solvate with non-halogenated organic solvent with the exception of water as a solvent.

10. A pharmaceutical composition, comprising azithromycin in the form of a solvate
with non-halogenated solvent with the exception of water as a solvent, or
azithromycin which is free from halogenated organic solvent in a stable
5 anhydrous form, in association with at least one pharmaceutical carrier or
diluent.

10



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 17/00, A61K 31/70	A3	(11) International Publication Number: WO 99/58541 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/EP99/03134 (22) International Filing Date: 6 May 1999 (06.05.99) (30) Priority Data: 9809939.3 8 May 1998 (08.05.98) GB 9812743.4 12 June 1998 (12.06.98) GB (71) Applicant (for all designated States except US): BIOCHEMIE S.A. [ES/ES]; Apartado de Correos, 97, E-08400 Granollers (ES). (72) Inventors; and (75) Inventors/Applicants (for US only): BOSCH, Immaculada [ES/ES]; Raval Cortines, 21, E-08500 Vic (ES). CENTEL-LAS, Victor [ES/ES]; Plaza Joan Asina, 8, 4-2-a, E-08440 Cardedeu (ES). DIAGO, José [ES/ES]; Calle Josep Carner, 43, 5-a, E-08400 Granollers (ES). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 20 January 2000 (20.01.00)
(54) Title: IMPROVEMENTS IN MACROLIDE PRODUCTION (57) Abstract <p>A process for the production of azithromycins by methylation of the nitrogen atom in position 9 of the ring structure in 9-deoxo-9a-aza-9a-homoerythromycins in the presence of formaldehyde and in the presence of a reducing agent in non-halogenated solvent; and azithromycin which is free from halogenated organic solvent in a stable anhydrous form; or in the form of a solvate with non-halogenated solvent with the exception of water.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

P.../EP 99/03134

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07H17/00 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP 0 109 253 A (PFIZER) 23 May 1984 (1984-05-23) page 7, line 4 - line 14 claims 1-3 ---	10 1,5,8,9
X A	EP 0 467 331 A (PLIVA PHARM & CHEM WORKS) 22 January 1992 (1992-01-22) page 3, line 17 - line 19 page 18, line 55 -page 19, line 2 claims 1,16 ---	10 1,5,8,9
X	EP 0 101 186 A (PFIZER) 22 February 1984 (1984-02-22) claims 9,11 page 6, line 26 -page 7, line 22 --- -/-	10

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

12 November 1999

Date of mailing of the international search report

26/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Held, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/03134

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 298 650 A (PFIZER) 11 January 1989 (1989-01-11) page 2, line 13 - line 54 claims ---	5-10
A	CHEMICAL ABSTRACTS, vol. 121, no. 23, 5 December 1994 (1994-12-05) Columbus, Ohio, US; abstract no. 280442, VAJTNER, Z. ET AL: "Kinetics of azithromycin synthesis from azaerythromycin by the HPLC method" XP002122554 abstract & KEM. IND. (1994), 43(6), 283-6 , ---	1,5,8,9
P,X	EP 0 879 823 A (HOVIONE INT LTD) 25 November 1998 (1998-11-25) column 1, line 24 -column 2, line 49 column 3, line 4 - line 10 claims 6,7 -----	1-5,8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/03134

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0109253 A	23-05-1984	AT 30237 T	15-10-1987
		AU 544790 B	13-06-1985
		AU 2130983 A	24-05-1984
		CA 1239639 A	26-07-1988
		CA 1250284 A	21-02-1989
		CS 8407926 A	16-07-1985
		CS 8308455 A	16-07-1985
		DD 216017 A	28-11-1984
		DE 3374065 A	19-11-1987
		DK 518683 A,B,	16-05-1984
		EG 16641 A	30-08-1990
		ES 527249 A	01-10-1985
		ES 543638 A	16-01-1986
		FI 834163 A,B,	16-05-1984
		GR 79425 A	22-10-1984
		IE 56234 B	22-05-1991
		IL 70228 A	31-03-1987
		JP 1040038 B	24-08-1989
		JP 1557268 C	16-05-1990
		JP 59104398 A	16-06-1984
		KR 8500968 B	02-07-1985
		NO 834146 A	16-05-1984
		NZ 206259 A	08-08-1986
		PH 19293 A	04-03-1986
		PH 21560 A	11-12-1987
		PT 77645 A,B	01-12-1983
		SU 1272996 A	23-11-1986
		US 4526889 A	02-07-1985
		YU 197285 A	31-08-1986
		YU 224383 A	30-04-1986
		ZA 8308460 A	26-06-1985
EP 0467331 A	22-01-1992	AT 129715 T	15-11-1995
		BG 60593 B	29-09-1995
		CA 2046956 A,C	19-01-1992
		CN 1059728 A,B	25-03-1992
		CS 9102195 A	18-03-1992
		DE 69114198 D	07-12-1995
		DE 69114198 T	27-06-1996
		DK 467331 T	05-02-1996
		ES 2081397 T	01-03-1996
		GR 3018330 T	31-03-1996
		HR 920491 A	31-08-1994
		HU 215158 B	28-12-1998
		HU 9500748 A	28-12-1995
		JP 2657132 B	24-09-1997
		JP 5132497 A	28-05-1993
		PL 166379 B	31-05-1995
		RO 109338 A	30-01-1995
		SI 9011409 A	31-10-1995
		SK 279075 B	03-06-1998
		RU 2045533 C	10-10-1995
		US 5250518 A	05-10-1993
EP 0101186 A	22-02-1984	US 4474768 A	02-10-1984
		AT 23343 T	15-11-1986
		AU 540056 B	01-11-1984
		AU 1692383 A	26-01-1984

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/03134

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0101186 A		CA 1202963 A	08-04-1986
		CA 1202619 C	01-04-1986
		CA 1202620 C	01-04-1986
		DK 332583 A,B,	20-01-1984
		FI 832606 A	20-01-1984
		FI 864479 A,B,	04-11-1986
		IE 55365 B	29-08-1990
		IL 69265 A	31-08-1986
		JP 1036834 B	02-08-1989
		JP 1554109 C	04-04-1990
		JP 59031794 A	20-02-1984
		JP 1193292 A	03-08-1989
		NO 832616 A	20-01-1984
		NZ 204938 A	11-04-1986
		PH 18054 A	18-03-1985
		PT 77062 A,B	01-08-1984
		ZA 8305204 A	27-02-1985
		CS 8305758 A	15-08-1985
		DD 215787 A	21-11-1984
		EG 16882 A	30-06-1989
		ES 527250 A	01-12-1985
		ES 544877 A	16-01-1986
		ES 544878 A	16-01-1986
		GR 77556 A	24-09-1984
		KR 8500963 B	02-07-1985
		SU 1274626 A	30-11-1986
		YU 133285 A	28-02-1986
		YU 207583 A	31-12-1985
EP 0298650 A	11-01-1989	WO 8900576 A	26-01-1989
		AP 44 A	27-07-1989
		AT 72446 T	15-02-1992
		AU 604553 B	20-12-1990
		BA 98213 B	02-08-1999
		BG 47348 A	15-06-1990
		CA 1314876 A	23-03-1993
		CN 1030422 A,B	18-01-1989
		CS 8804896 A	14-03-1990
		CY 1776 A	20-10-1995
		DD 271705 A	13-09-1989
		DE 3868296 A	19-03-1992
		DK 380688 A	10-01-1989
		FI 900087 A,B,	08-01-1990
		GR 3003737 T	16-03-1993
		HK 127594 A	25-11-1994
		HU 9500738 A	28-11-1995
		IE 60354 B	29-06-1994
		IL 86979 A	15-11-1992
		IN 168879 A	29-06-1991
		JP 1038096 A	08-02-1989
		JP 1903527 C	08-02-1995
		JP 6031300 B	27-04-1994
		LV 10624 A	20-04-1995
		MX 12213 A	01-05-1993
		NZ 225338 A	26-02-1990
		OA 8743 A	31-03-1989
		PT 87933 A,B	30-06-1989
		RO 107257 A	30-10-1993

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

P.../EP 99/03134

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0298650	A	SG 27794 G	14-10-1994
		SI 8811325 A	31-12-1996
		RU 2066324 C	10-09-1996
		YU 132588 A	28-02-1990
EP 0879823	A	PT 102006 A	31-12-1998
	25-11-1998	AU 6193898 A	19-11-1998
		CA 2237747 A	19-11-1998
		CN 1199736 A	25-11-1998
		HU 9801094 A	28-12-1998
		JP 10316699 A	02-12-1998
		NZ 330228 A	29-06-1999